



teach with space

→ CLOUD CHAMBER

Radioactivity in a cosmic setting





Fast facts page 3

Background page 4

Experiment preparation page 6

Answers to discussion questions page 8

Answers to student worksheets page 11

Links page 12

→ CLOUD CHAMBER

Radioactivity in a cosmic setting

FAST FACTS

Age range: 14-18 years old

Type: teacher demonstration/group practical

Complexity: medium

Teacher preparation time: 5 minutes per cloud chamber

Lesson time required: 1 hour

Cost per kit: medium (5-25 euros per cloud chamber)


Location: small indoor space

Includes use of: low level radiation source, dry ice (solid CO₂), alcohol

Students will learn

- Fundamental particles
- Particle interactions
- Atomic number, mass number
- Radiation and radioactive decay
- Radioactive decay as a power source for spacecraft
- The negative effects of ionising radiation on electronics
- Interactions between charged particles and magnetic fields
- Cosmic rays including their interactions with the atmosphere

You also need



↑ [Cloud chamber video \(VP03\)](#). See [Links](#) section.

Outline

Cloud chambers are boxes specially made to detect charged particles and radiation. In this activity, a cloud chamber is used to observe alpha and beta particles, the charged products of the radioactive decay of thorium-232. Students should already have been introduced to the concept of radioactive decay and the differences between alpha and beta radiation. This experiment will assist students in their understanding by providing a physical demonstration of radioactive decay.

Students will observe the condensation trails that charged particles leave in their wake in a cloud chamber, and learn to identify particles based on the trail properties. Trails made by cosmic rays can be seen in the cloud chamber, which could provide a starting point for introducing cosmic rays and their deflection by the Earth's magnetic field. Students will practice calculating and writing radioactive decay equations. Implications of radiation for ESA are discussed, including the negative effects of radiation on circuits, and the use of radioactive decay as a power source.

Students will improve

- General experimental skills including using equipment appropriately and making and recording observations.
- The communication and discussion of results, through asking relevant questions to expand understanding and knowledge of a subject.
- Their application of knowledge gained through experimental observations to solve theoretical problems.

→ BACKGROUND

With the invention of the first cloud chamber, a new branch of physics was born: the study of particles and their interactions. Cloud chambers allow us to observe charged particles to determine their properties so we can understand them in more detail.

Inside the chamber, alcohol molecules in vapour form are ionised when charged particles pass close to them. This causes the alcohol vapour to condense into liquid droplets, resulting in a visible condensation trail (contrail) where the charged particle interacted with the alcohol. The length and thickness of this trail depends on the mass of the particle.

These charged, ionising particles come from a range of sources including materials undergoing radioactive decay. During radioactive decay, unstable atoms break up into smaller, more stable atoms, releasing either alpha radiation (a particle made up of two neutrons and two protons) or beta radiation (an electron or a **positron***).




Figure 1

↑ ESA's Swarm satellites orbit the Earth, studying its magnetic field.




Figure 2

↑ Earth's magnetic field. Note that the rotation axis and the magnetic field axis are aligned differently and they do not intersect at the centre of the Earth

As well as arising from radioactive decay, a lot of ionising radiation is present in space, originating from sources such as **cosmic rays***, the **solar wind***, and **coronal mass ejections***. Moving charged particles are deflected by magnetic fields, so the Earth is largely shielded from ionising space radiation by its own magnetic field, the magnetosphere, which is currently being studied by ESA's Swarm (Figure 1) and Cluster satellites. The magnetosphere is generated by the motion of the magnetic elements iron and nickel in the Earth's core. The magnetic field lines of the Earth are shown in Figure 2.

* **Positron**: a particle with the same mass as an electron, but with the opposite charge.


Cosmic rays: very high-energy particles, mostly protons and atomic nuclei, originating from the Sun and objects outside the Solar System.

Solar wind: a stream of charged particles emitted from the Sun – mainly electrons and protons.

Coronal mass ejection: a burst of electromagnetic radiation, particles, and magnetic fields ejected from the solar corona into space.

Charged particles and radiation can get trapped within the Earth's magnetic field lines which are concentrated in the inner and outer **Van Allen radiation belts*** depicted in Figure 3. This can have an adverse effect on electronic equipment on board spacecraft orbiting the Earth. However, radiation can also be used for good. Spacecraft can be powered via the Seeback effect; a temperature difference between two electrical conductors (induced by heating one with radiation) can produce a voltage difference between the two, causing current to flow.

A cloud chamber can be developed further by including a magnetic field, which deflects moving charged particles. The direction and amount of deflection depends on the charge of the particle. This fact was instrumental in the discovery of antimatter in 1932 when Carl Anderson used a cloud chamber with a magnetic field to observe cosmic rays, charged particles from space, and found that some of the particles were deflected in the opposite direction to what was expected of them.



↑ Van Allen radiation belts. Two doughnut-shaped rings that circle the Earth and contain trapped charged particles.

* **Van Allen radiation belts**: layers of energetically charged particles held in place around the Earth by the magnetic field. The altitude ranges from about 1000 to 60000 km above the Earth's surface.

→ EXPERIMENT PREPARATION

The teach with space - cloud chamber | VPO3 video, available on ESA's Education website, shows the complete set-up and use of the cloud chamber. This video can be used to guide students on setting up their chambers, to explain what they are observing, or simply as an alternative to building cloud chambers in the classroom.

Equipment for each experiment

To prepare in advance

- Medium-sized plastic fish tank
- Adhesive-backed felt (or normal felt and glue appropriate for felt and plastic)

To perform the experiment

- Two thoriated tungsten welding rods (or other alpha/beta source)
- About 2.5 kg dry ice (solid CO₂)
- 20 ml isopropyl alcohol, also known as isopropanol (or ethanol if this is unavailable)
- A plastic fish tank with pre-attached felt
- Two metal trays (baking trays work well)
- One piece of black card or laminated black paper (to line trays if they are not dark)
- One or two intense light sources (e.g. an LED light strip, torch, or slide projector light)
- A sheet of paper to wrap around one rod
- Poster putty or reusable adhesive
- A pipette
- Thermal protective gloves
- Safety glasses (one pair per person)

Set-up

1. Cut strips of felt to a width of approximately 4 cm and the length of the tank sides.
2. With the fish tank open face up, line each of the four sides of the fish tank with the felt strips (they should be near the top of the chamber when in use – see figure on student worksheet). Felt that comes with an adhesive on the back is easiest, but glue can be used to attach the felt. The alcohol will act as a solvent, so any adhesives should be tested before use.
3. Instructions on how to carry out the experiment are provided in the student activities document (teach with space - cloud chamber | P03b).

Health & Safety

Refer to your school's policy and the national safety guidelines for your country and complete a risk assessment.

Isopropanol (and other alcohol if used):

- Isopropanol is highly flammable and an irritant. Do not use near fire and wear goggles.
- Ensure the room is well ventilated and the lid is immediately replaced on the propanol bottle.

Dry Ice:

- Ensure the room is well ventilated to avoid high carbon dioxide levels.
- Extensive inhalation of cold vapours can result in lung damage/asthma attacks.
- Wear eye protection and thermally insulated gloves to avoid cold burns from direct contact with the dry ice or cold equipment. If a dry ice burn is obtained, treat as a normal burn - soak under cold water for 10 minutes and seek further medical assistance if necessary.
- Dry ice should NOT be stored in a fully sealed container, as carbon dioxide gas will build up. The container should be clearly labelled.

Thoriated tungsten rods – radiation source:

- The thoriated tungsten rods contain thorium oxide, an emitter of alpha particles. It has a very low penetration and therefore is safe to handle when taking appropriate precautions.
- It is recommended that large numbers of rods are stored in a metal box with a radioactive warning on, and stored along with other radioactive sources in your school.
- If the rod is showing signs of flaking or disintegrating it should be disposed of appropriately.
- No part of the rod should be ingested. Whilst this is very unlikely, in the event of ingestion of any part of the rod, seek medical attention before following your school's radioactive substances policy.

→ ANSWERS TO DISCUSSION QUESTIONS

1. Where do most of the contrails seem to originate?

They mostly originate from the two thoriated tungsten rods.

2. We are observing the radioactive decay of thoriated rods. What could the contrails be showing?

The contrails are showing the paths taken by alpha and beta particles produced in the radioactive decay of thorium-232 in the welding rods.

3. Are there differences between contrails from each of the thoriated rods?

From the unwrapped rod, both long thin tracks and short thick tracks can be seen. From the wrapped rod, only long thin tracks can be seen.

4. What is the difference between alpha and beta decay?

Radioactive decay is the process by which the nucleus of an unstable atom loses energy by emitting ionising radiation in order to correct the ratio of protons to neutrons in its nucleus. There are two main types of radioactive decay: alpha and beta.

In alpha decay, two protons and two neutrons are emitted. This leaves a daughter particle with a mass number that is four less and an atomic number that is two less than the original atom. For example, uranium-238 decays by alpha emission, losing two protons and two neutrons, to become thorium-234.

For beta decay, there are two options:

β^- : A neutron transforms into a proton, emitting an electron in the process.

β^+ : A proton transforms into a neutron, emitting a positron in the process.

5. Could the differences between alpha and beta decay have any effect on the contrails that you see?

Alpha particles are physically bigger and more ionising than beta particles because they have a charge of +2 from the two protons. They have a relatively short journey before all of their kinetic energy is transferred. This means that contrails from alpha particles are the shorter, thicker trails, about 2-5 cm in length. As they have a weaker charge, beta particles are less ionising and so they travel further before losing all of their energy. They produce longer trails of up to 10 cm.

Only beta particles are observed from the rod wrapped in paper. This is because alpha particles give away all of their energy to the paper particles by ionising them, before they can reach the alcohol vapour. Beta particles are less strongly ionising and so they travel straight through the paper surrounding the rod without interacting with it. The very short penetrating distance of alpha radiation is clear – alpha particles lose all of their energy in a short distance.



↑ A thoriated rod emitting alpha and beta radiation.

6. How are contrails formed in a cloud chamber?

Alcohol in the felt is very volatile and quickly forms a vapour at the top of the cloud chamber. As the bottom of the cloud chamber reaches a low temperature due to the dry ice, the alcohol vapour cools rapidly as it falls. The alcohol would naturally condense into a liquid at this lower temperature, as it is below its boiling point.

However, in order to condense, particles of gas need a 'seed', or a small particle or disturbance on which they can grow. As there are no seeds in the chamber, a thin 'super-saturated' layer of alcohol that is unable to condense gathers around the bottom of the cloud chamber.

A charged particle (such as an alpha particle produced through radioactive decay) passing through the chamber ionises alcohol molecules in its path. These ionised alcohol molecules act as a seed for the surrounding neutral alcohol molecules, allowing them to condense into droplets, which form the contrail that can be seen in the chamber. Soon after the droplets are formed, they fall to the bottom of the cloud chamber and disperse; the contrail is only visible for a couple of seconds.

7. Cosmic rays are high energy particles from space. How do we observe them?


Occasionally you may see contrails that do not appear to originate from the thoriated tungsten rods. While these can be the products of random radioactive decays outside of the chamber, they are more commonly the result of cosmic rays from the Sun and other objects in the Universe such as stars and galaxies. Cosmic rays from space enter the atmosphere and interact with atmospheric particles. These interactions firstly create particles called **pions***. Pions decay very quickly, often into **muons***. These muons can be observed as extremely long, thin, straight trails in the cloud chamber.

8. What could be the effects of radiation on spacecraft?

Ionising radiation can knock electrons loose within electric circuits, leading to interference with electrical systems on Earth-orbiting satellites. In most cases this effect is simply observed as a reduction in the quality of the data, but effects are magnified during events such as coronal mass ejections due to an increased intensity of ionising particles. This can lead to excess current through a circuit, which can result in damage or destruction of equipment. As a result, satellite and spacecraft engineers have to protect delicate circuits from harmful space radiation. The simplest approach to this is to shield them with specially designed material containing atoms with a high atomic number, and therefore many protons and neutrons that can absorb much of the space radiation.

9. How can radioactive decay be used as a power source in space?

Radioactive decay produces energy, which causes material to heat up. The Seebeck effect makes use of this; it is a phenomenon in which a temperature difference between two different electrical conductors can produce a voltage difference between the two. One conductor is heated by radioactive decay, whilst the other conductor is kept cool, for example by exposing it to the cold space environment. Heated electrons will flow from the hotter conductor to the cooler conductor. If the conductors are connected through an electrical circuit, direct current will flow within the circuit (Figure 5). This method of power production has been used in many spacecraft, like ESA's Ulysses mission and the ESA/NASA/ASI Cassini-Huygens mission.



↑ The Seebeck effect.

* **Pion (Pi meson)**: a type of particle much smaller than an atom. There are three different pions; π^0 , π^+ , and π^- .
* **Muon**: a particle with the same charge as an electron but with much more mass.

Did you know?


ESA's Cluster mission studies how the constant stream of charged atoms and electrons from the Sun affects the Earth's magnetosphere. This stream of particles is called the solar wind and gusts of solar wind have the potential to affect power supplies here on Earth (a strong gust could stop all electronic devices from working!). Solar wind can also harm the electronics on-board satellites. Cluster studies phenomena that can cause damage in this way - it assists us in preparing for sudden, potentially dangerous, bursts of solar energy!




→ ANSWERS TO STUDENT ACTIVITIES

1. Decay reactions


a. α decay




b. β^- decay (electron emission):



c. β^+ decay (positron emission):



2. Thorium decay



3. The stable element produced at the end of the decay chain is **lead**.

Links

Teach with space collection

ESA teach with space - cooking a comet video | VP03: www.esa.int/spaceinvideos/Videos/2014/07/Cloud_chamber_classroom_demonstration_video_VP03

ESA related missions and science

ESA Cassini-Huygens mission: www.esa.int/Our_Activities/Our_Activities/Space_Science/Cassini-Huygens

Cassini's solstice mission: saturn.jpl.nasa.gov/

ESA Ulysses mission: www.esa.int/Our_Activities/Space_Science/Ulysses_overview

ESA Swarm mission: www.esa.int/Our_Activities/Observing_the_Earth/The_Living_Planet_Programme/Earth_Explorers/Swarm

ESA Cluster mission: www.esa.int/Our_Activities/Space_Science/Cluster/Science_objectives

ESA radioactive decay articles

Radioactive decay of titanium powers supernova remnant: www.esa.int/Our_Activities/Space_Science/Radioactive_decay_of_titanium_powers_supernova_remnant

Radioactive iron: www.esa.int/Our_Activities/Space_Science/Integral/Radioactive_iron_a_window_to_the_stars

teach with space – cloud chamber | P03a

www.esa.int/education

The ESA Education Office welcomes feedback and comments
teachers@esa.int

An ESA Education production

Copyright © European Space Agency 2016